arXiv:1403.7826v1 [math.DS] 30 Mar 2014

PURE DISCRETE SPECTRUM FOR A CLASS OF ONE-DIMENSIONAL SUBSTITUTION TILING SYSTEMS

MARCY BARGE

Abstract. We prove that if a primitive and non-periodic substitution is injective on initial letters, constant on final letters, and has Pisot inflation, then the $\mathbb{R}$-action on the corresponding tiling space has pure discrete spectrum. As a consequence, all β-substitutions for β a Pisot simple Parry number have tiling dynamical systems with pure discrete spectrum, as do the Pisot systems arising, for example, from the Jacobi-Perron and Brun continued fraction expansions.

Contents

1. Introduction 2

2. Background and Notation 6

3. Proof of the Main Theorem 9

4. Applications 28

4.1. β-substitutions 28

Date: April 1, 2014.

2010 *Mathematics Subject Classification.* Primary: 37B05, 37B50 ; Secondary: 11A55, 11A63 .

Key words and phrases. Substitution, Tiling Space, Discrete Spectrum, Maximal Equicontinuous Factor.

4.2. Arnoux-Rauzy, Brun, and Jacobi-Perron substitutions 30

References 32

1. Introduction

Substitution dynamical systems arise in materials science (when analyzing spectral properties of atomic arrangements), in number theory (when investigating arithmetical properties of expansions in various bases), in hyperbolic dynamics (in the construction of Markov partitions for toral automorphisms), in computer science, word combinatorics, and, generally, when considering hierarchical, self-similar structures. A natural and effective approach to understanding such a system is to compare it with an action by translation on a compact abelian group: in the 'best-behaved' case, when the system is measurably isomorphic with such an action, the system is said to have *pure discrete spectrum*.

There are two dynamical systems commonly associated with a substitution ϕ: a subshift denoted by $(\Sigma_\phi, \mathbb{Z})$ called the *substitutive system*; and the *tillng dynamical system* $(\Omega_\phi, \mathbb{R})$, which is a particular suspension of the substitutive system. Pure discreteness of the substitutive system implies pure discreteness of the tillng dynamical system, but the converse does not hold without additional conditions (see [CS]). In this article we deal exclusively with the tiling dynamical system - it has the important advantage of

supporting a substitution induced homeomorphism Φ which interacts with the translation $\mathbb{R}$-action via $\Phi(T-t) = \Phi(T) - \lambda t$ for tilings $T \in \Omega_\phi$ and real numbers t. Here $\lambda > 1$, the *inflation* associated with ϕ, is the largest eigenvalue of the *abelianization*, or *substitution matrix*, A of ϕ whose ij-th entry is the number of times the i-th letter appears in the image under ϕ of the j-th letter. In order for the tiling dynamical system $(\Omega_\phi, \mathbb{R})$ to have pure discrete spectrum it is necessary that the inflation λ be a Pisot number (an algebraic integer, all of whose other algebraic conjugates lie strictly inside the unit circle) (see [S1]). A substitution with Pisot inflation is called a *Pisot substitution* and a substitution is said to be *irreducible* if the characteristic polynomial of its substitution matrix is irreducible over $\mathbb{Q}$. We will denote the alphabet of a substitution by $\mathcal{A}$.

There are various algorithms for determining whether or not a given substitution dynamical system has pure discrete spectrum (see, in particular, [AL1, AL2]); these may involve a considerable amount of computation and it is desirable to have, instead, simple and easily checked conditions that guarantee pure discrete spectrum. It is conjectured, for example, that the system associated with an irreducible Pisot substitution has pure discrete spectrum (this is known as the *Pisot substitution conjecture*, see [ABBLS]), but this has only been verified for substitutions on two letters ([HS]). It happens that many substitutions ϕ that arise in applications have the following special form: every

letter of the alphabet $\mathcal{A}$ occurs as a first letter of a word $\phi(a)$ for some $a \in \mathcal{A}$ and all such words end in the same letter. It is the main theorem of this article that the tiling dynamical system of any (primitive) Pisot substitution with this property has pure discrete spectrum.

Given a substitution ϕ on the finite alphabet $\mathcal{A}$, say $\phi(a) = a_1 \cdots a_{t(a)}$ for $a \in \mathcal{A}$, we will say that ϕ is *injective on initial letters* if the map $a \mapsto a_1$ is injective, and we will say that ϕ is *constant on final letters* if $a_{t(a)} = b_{t(b)}$ for all $a, b \in \mathcal{A}$. Examples of families of substitutions that are both injective on initial letters and constant on final letters include β-substitutions for β a simple Parry number, and substitutions arising in connection with continued fraction algorithms (see Section 4). The following theorem is the main result of this article.

Theorem (Theorem 3.12 in text): If ϕ is a primitive Pisot substitution that is injective on initial letters and constant on final letters then the tiling dynamical system $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum.

In outline, the argument will be as follows:

(1) Every tiling dynamical system $(\Omega, \mathbb{R})$ has a *maximal equicontinuous factor* $(\Omega_{\max}, \mathbb{R})$ and $(\Omega, \mathbb{R})$ has pure discrete spectrum if and only if the factor map $\pi_{max} : \Omega \to \Omega_{max}$ is a.e. one-to-one.

(2) Given $T, T' \in \Omega_\phi$ we write $T \approx_s T'$ provided $\pi_{max}(T) = \pi_{max}(T')$ and, for a dense set of $t \in \mathbb{R}$, there is $n(t) \in \mathbb{N}$ so that so that $\Phi^{n(t)}(T-t)$ and $\Phi^{n(t)}(T'-t)$ have exactly the same tiles at the origin. The relation $\approx_s$ is a closed equivalence relation.

(3) If the system $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum, the quotient system $(\Omega_\phi / \approx_s, \mathbb{R})$ is isomorphic with a tiling dynamical system $(\Omega_{\phi_s}, \mathbb{R})$ for a primitive, non-periodic Pisot substitution ϕ_s that is also injective on initial letters and constant on final letters. The relation $\approx_s$ is trivial on Ω_{ϕ_s}.

(4) If ψ is a primitive, non-periodic, Pisot substitution that is injective on initial letters and constant on final letters, then $\approx_s$ is nontrivial on Ω_ψ.

The proof now follows by contradiction: Were $(\Omega_\phi, \mathbb{R})$ not to have pure discrete spectrum, the relation $\approx_s$ would be trivial on Ω_{ϕ_s}. But item (4) would apply with $\psi = \phi_s$ to say that $\approx_s$ is nontrivial on Ω_{ϕ_s}.

Many of the ingredients of the proof have been developed elsewhere: (1) holds quite generally and is a consequence of the Halmos - von Neumann theory (see Chapter 3

of [W], for example); that Φ is a homeomorphism is a result of Mossé ([M], or, more generally, [S2]); much of (2) and (3) appears in [B] in arbitrary dimension; and (4) is derived from the main idea of [BD1]. The details are presented in Section 3 and applications are given in Section 4.

2. Background and Notation

Given an alphabet $\mathcal{A}$, $\mathcal{A}^*$ will denote the set of all finite, nonempty, words in $\mathcal{A}$. A substitution $\phi : \mathcal{A} = \{1, \dots, d\} \to \mathcal{A}^*$, $d > 1$, is *primitive* if some power of its abelianization A is strictly positive. In this case A has a unique (up to scale) positive left eigenvector $\omega = (\omega_1, \dots, \omega_d)$. The corresponding Perron-Frobenius eigenvalue λ is called the *dilation* of ϕ. The *prototiles* for ϕ are the labeled intervals $\rho_i := ([0, \omega_i], i)$. (For convenience, we will sometimes confuse ρ_i with $[0, \omega_i] \times \{i\}$.) A *tile* is a translate of a prototile, $\rho_i - t := ([-t, \omega_i - t], i)$, with *support* $spt(\rho_i - t) := [-t, \omega_i - t]$ and *type* i. A *tiling* is a collection T of tiles whose supports cover $\mathbb{R}$ with the property that any two distinct tiles in T have supports with disjoint interiors. A *patch* P is a finite subset of a tiling with *support* equal to the union of the supports of its constituent tiles. Given a tile $\tau = \rho_i - t$, let $\Phi(\tau)$ be the patch

$$\Phi(\tau) = \{\tau_1, \dots, \tau_k\},$$

with

$$\tau_j := \rho_{i_j} - \lambda t + \sum_{l=1}^{j-1} \omega_{i_l},$$

where $\phi(i) = i_1 \cdots i_k$. Extend Φ to patches P by $\Phi(P) := \cup_{\tau \in P} \Phi(\tau)$, and likewise to tilings. Notice that $spt(\Phi(P)) = \lambda \cdot spt(P)$.

Given a primitive substitution ϕ as above, there are $k \in \mathbb{N}$ and $a, b \in \mathcal{A}$ so that $\phi^k(a) = a \cdots$, $\phi^k(b) = \cdots b$, and ba is in the *language*, $\mathcal{L} := \{w : w \text{ is a factor of } \phi^n(i) \text{ for some } n \in \mathbb{N} \text{ and } i \in \mathcal{A}\}$, of ϕ. Let $P = \{\rho_b - \omega_b, \rho_a\}$ and let $T = \cup_{m \in \mathbb{N}} \Phi^{mk}(P)$. Then T is a tiling and $\Phi^k(T) = T$. The *tiling space* of ϕ, or *hull* of T, is

$$\Omega_\phi := cl\{T - t : t \in \mathbb{R}\},$$

with the closure taken in the *local topology* in which two tilings are close if a small translate of one agrees with the other in a large neighborhood of the origin. This is a metric topology (we will denote the metric by d), in which Ω_ϕ is compact and connected. Furthermore, the space Ω_ϕ does not depend on T and the $\mathbb{R}$-action by translation, $(S, t) \mapsto S - t$, on Ω_ϕ is minimal and uniquely ergodic (see, for example, [AP]).

For ϕ that is primitive and non-periodic (that is, there are no translation periodic tilings in Ω_ϕ), the map $\Phi : \Omega_\phi \to \Omega_\phi$ is a homeomorphism ([M],[S2]) satisfying $\Phi(T -$

$t) = \Phi(T) - \lambda t$ for all $T \in \Omega_\phi$, $t \in \mathbb{R}$.

Given $T \in \Omega_\phi$ and $R \geq 0$, the *R-patch of* T *at 0* is defined as

$$B_R[T] := \{\tau \in T : spt(\tau) \cap \bar{B}_R(0) \neq \emptyset\}.$$

(Here $\bar{B}_R(0)$ is the closed ball of radius R at the origin.) Thus $d(T, T')$ is small if there is a t with $|t|$ small, and a large R, so that $B_R[T] = B_R[T' - t]$.

The substitution ϕ is a *Pisot substitution* if its inflation λ is a Pisot number. For such ϕ, the maximal equicontinuous factor $(\Omega_{\max}, \mathbb{R})$ of the tiling dynamical system $(\Omega_\phi, \mathbb{R})$, which is unique up to topological isomorphism, is a (non-trivial) torus or solenoid of dimension equal to the algebraic degree of λ, the factor map $\pi_{max} : \Omega_\phi \to \Omega_{max}$ is uniformly finite-to-one, measure preserving (with respect to the unique invariant measure on Ω_ϕ and Haar measure on Ω_{max}), and almost everywhere r-to-one for an $r < \infty$ called the *coincidence rank* of ϕ ([BKw, BBK]). The system $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum if and only if the coincidence rank, r, equals 1. Besides semi-conjugating $\mathbb{R}$-actions, π_{max} also semi-conjugates $\mathbb{Z}$-actions: there is a hyperbolic and algebraic automorphism $\Phi_{max} : \Omega_{max} \to \Omega_{max}$ with $\pi_{max} \circ \Phi = \Phi_{max} \circ \pi_{max}$.

Tilings T and T' are *proximal* if $inf_{t \in \mathbb{R}} d(T - t, T' - t) = 0$ and *strongly proximal* if for all $R > 0$ there is $t_R \in \mathbb{R}$ with $B_0[T - t_R] = B_0[T' - t_R]$. These two relations are equal for Pisot ϕ (this is a consequence of the 'Meyer property' - see [BK]) and clearly $\pi_{max}(T) = \pi_{max}(T')$ if T and T' are proximal. The equicontinuous structure relation for

Pisot ϕ is *strong regional proximality*, $\sim_{srp}$, defined by: $T \sim_{srp} T'$ if and only if for each $R > 0$ there are $S_R, S'_R \in \Omega_\phi$ and $t_R \in \mathbb{R}$ so that $B_R[T] = B_R[S]$, $B_R[T'] = B_R[S'_R]$, and $B_R[S_R - t_R] = B_R[S'_R - t_R]$. That is, $\pi_{max}(T) = \pi_{max}(T')$ if and only if $T \sim_{srp} T'$ (see [BK]).

3. Proof of the Main Theorem

Suppose that ϕ is a (primitive and non-periodic) Pisot substitution on the alphabet $\mathcal{A} = \{1, \ldots, d\}$. Tilings $T, T' \in \Omega_\phi$ are *eventually coincident at* $t \in \mathbb{R}$ if there is $n \in \mathbb{N}$ so that $B_0[\Phi^n(T - t)] = B_0[\Phi^n(T' - t)]$. Note that if T, T' are eventually coincident at t, then there is $\epsilon > 0$ so that T, T' are eventually coincident at t' for all $t' \in (t - \epsilon, t + \epsilon)$. We will say that T and T' are *densely eventually coincident* if T, T' are eventually coincident at t for a set of t dense in $\mathbb{R}$ (hence for an open dense set of t). Define $\approx_s$ on Ω_ϕ by $T \approx_s T'$ if and only if T and T' are strongly regionally proximal and densely eventually coincident.

The following theorem is a compilation of one-dimensional restrictions of several results of [B]. We include a proof for completeness.

Theorem 3.1. *$\approx_s$ is a closed equivalence relation, invariant under translation and under Φ, and $(\Omega_\phi/\approx_s, \mathbb{R})$ is the maximal equicontinuous factor of $(\Omega_\phi, \mathbb{R})$ if and only if $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum. The maximal equicontinuos factor map on Ω_ϕ*

factors through the quotient map from Ω_ϕ to $\Omega_\phi/\approx_s$ and that quotient map is a.e. one-to-one. In particular, if $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum, then neither does $(\Omega_\phi/\approx_s, \mathbb{R})$.

Proof. As mentioned in the previous section, $T \sim_{srp} T'$ is equivalent to $\pi_{max}(T) = \pi_{max}(T')$, so $\sim_{srp}$ is a closed equivalence relation. That 'T is densely eventually coincident with T'' really means 'T is open-and-densely eventually coincident with T'' implies that dense eventual coincidence is an equivalence relation, so $\approx_s$ is an equivalence relation. To see that it's closed, suppose that $T_n \to T$, $T'_n \to T'$ and $T_n \approx_s T'_n$ for each n. Then $T \sim_{srp} T'$. Fix $R > 0$ and let $t_n, t'_n \to 0$ be so that (for large n) $B_R[T - t_n] = B_R[T_n]$ and $B_R[T' - t'_n] = B_R[T']$. By Corollary 5.8 of [BK] there are, up to translation, only finitely many pairs of the form $(B_R[S], B_R[S'])$ with $S \sim_{srp} S'$. It follows that $t_n = t'_n$ for infinitely many n. Since, for $|t| < R$, eventual coincidence of T_n and T'_n at t depends only on the R-patches of T_n and T'_n at the origin, and T_n, T'_n are densely eventually coincident, we see that T and T' are eventually coincident at a set of t dense in $B_R(0)$. Thus $T \approx_s T'$ and $\approx_s$ is a closed equivalence relation.

The Φ- and $\mathbb{R}$-invariance of $\approx_s$ is clear. Also, since the relation $\approx_s$ is contained in the relation $\sim_{srp}$, π_{max} factors through $\Omega_\phi/\approx_s$.

Suppose now that $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum. We will argue that $(\Omega_\phi/\approx_s, \mathbb{R})$

is the maximal equicontinuous factor of $(\Omega_\phi, \mathbb{R})$ by showing that strong regional proximality implies dense eventual coincidence. Suppose that $T \sim_{srp} T'$ and suppose that there are $t_0 \in \mathbb{R}$ and $\epsilon > 0$ so that T and T' are not eventually coincident at any $t \in (t_0 - \epsilon, t_0 + \epsilon)$. Pick $n_i \to \infty$ with $\Phi^{n_i}(T - t_0) \to S \in \Omega_\phi$ and $\Phi^{n_i}(T' - t_0) \to S' \in \Omega_\phi$. Then $S \sim_{srp} S'$. Suppose that $B_0[S-t] = B_0[S'-t]$ for some $t \in \mathbb{R}$. There are then (for large i) $t_i \to t$ with $B_0[\Phi^{n_i}(T-t_0)-t_i] = B_0[S-t] = B_0[S'-t] = B_0[\Phi^{n_i}(T'-t_0)-t_i]$ (we have again used that there are, up to translation, only finitely many pairs $(B_0[V], B_0[V'])$ with $V \sim_{srp} V'$). Then $B_0[\Phi^{n_i}(T - t_0 - (t_i/\lambda^{n_i}))] = B_0[\Phi^{n_i}(T' - t_0 - (t_i/\lambda^{n_i}))]$ so that T and T' are eventually coincident at $t_0 + (t_i/\lambda^{n_i})$. We have a contradiction when i is large enough so that $t_i/\lambda^{n_i} < \epsilon$. Thus $B_0[S - t] \neq B_0[S' - t]$ for all $t \in \mathbb{R}$. As there are only finitely many pairs $(B_0[S - t], B_0[S' - t])$, up to translation, we see that $d(S - t, S' - t)$ is bounded away from 0. Then by minimality of the $\mathbb{R}$-action and closed-ness of $\sim_{srp}$, $\pi_{max} : \Omega_\phi \to \Omega_{max}$ is at least 2-to-1 everywhere. But π_{max} is a.e. 1-to-1 since $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum. Hence T must be densely eventually coincident with T'. This proves that if $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum then $\sim_{srp}$ is the same as $\approx_s$; that is, $(\Omega_\phi/\approx_s, \mathbb{R})$ is the maximal equicontinuous factor.

Conversely, suppose that $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum. Then the coincidence rank of ϕ is $r \geq 2$ and there are $z \in \Omega_{max}$ and $T_1, \ldots, T_r \in \pi_{max}^{-1}(z)$ with T_i and T_j disjoint for $i \neq j$ (this is from [BK] - see Theorem 4 of [B]). By minimality (and the

fact that, up to translation, there are only finitely many pairs $(B_0[T_i - t], B_0[T_j - t]))$, this is true for all $z \in \Omega_{max}$. Such T_i and T_j are not proximal, and therefore $\Phi^n(T_i)$ and $\Phi^n(T_j)$ are not proximal for any $n \in \mathbb{N}$ (this is because Φ is a homeomorphism with $\Phi(S - t) = \Phi(S) - \lambda t$, so Φ^n preserves proximality for all $n \in \mathbb{Z}$). Then by Lemma 5.12 of [BK], $\Phi^n(T_i)$ and $\Phi^n(T_j)$ are disjoint for all $n \in \mathbb{N}$ and $i \neq j$. It follows that the factor map from $\Omega_\phi / \approx_s$ to Ω_{max} is at least r-to-1 everywhere. Since π_{max} is almost everywhere r-to-1, the quotient map from Ω_ϕ to $\Omega_\phi / \approx_s$ must be almost a.e. 1-to-1 (and, furthermore, $(\Omega_\phi / \approx_s, \mathbb{R})$ is not the maximal equicontinuous factor). □

Assume now that $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum. We will push the relation $\approx_s$ down to the (disjoint) union $\cup_{i=1}^d \rho_i$ of the prototiles for ϕ. Let the relation R on $\cup_{i=1}^d \rho_i$ be defined by $(x, i)R(y, j)$ if and only if there are $T, T' \in \Omega_\phi$ and $t, t' \in \mathbb{R}$ so that: $T \approx_s T'$, $t + \rho_i \in T$, $t' + \rho_j \in T'$, and $t + x = t' + y$.

Lemma 3.2. *: Suppose that ϕ is Pisot and is constant on final letters. There is then $\epsilon > 0$ so that if $(x_n, i_n)R(x_{n+1}, i_{n+1}]$ for $n = 1, \dots N - 1$ and P_1 and P_N are the patches $P_1 = \{\rho_1 - \omega_1 - x_1, \rho_{i_1} - x_1\}$ and $P_N = \{\rho_1 - \omega_1 - x_N, \rho_{i_N} - x_N\}$, then $[-\epsilon, 0] \subset spt(P_1) \cap spt(P_N)$ and P_1 and P_N are densely eventually coincident at s for a dense set of $s \in [-\epsilon, 0]$.*

Proof. Suppose $l \in \mathcal{A} = \{1, \dots, d\}$ is such that $\phi(a) = \cdots l$ for all $a \in \mathcal{A}$. Then the tiles $\rho_i - \omega_i$ and $\rho_j - \omega_j$ are eventually coincident at s for all $s \in (-\omega_l/\lambda, 0)$ for each $i, j \in \mathcal{A}$.

Let $\epsilon = \omega_l/\lambda$. The conclusion of the lemma is certainly true if $N = 1$. Suppose it to be true for some $N \geq 1$ and suppose that $(x_n, i_n)R(x_{n+1}, i_{n+1})$ for $n = 1, \ldots N$. Then P_1 and P_N are densely eventually coincident on $[-\epsilon, 0]$ and there are $T, T' \in \Omega_\phi$ and $t, t' \in \mathbb{R}$ so that: $T \approx_s T'$, $t+\rho_{i_N} \in T$, $t'+\rho_{i_{N+1}} \in T'$, and $t+x_N = t'+x_{N+1}$. Let $\rho_i-\omega_i+t$ and $\rho_j-\omega_j+t'$ be the tiles of T and T' immediately to the left of $t+\rho_{i_N}$ and $t'+\rho_{i_{N+1}}$. Then the patches $P'_N := \{\rho_i-\omega_i+t, t+\rho_{i_N}\}-t-x_N = \{\rho_i-\omega_i-x_N, \rho_{i_N}-x_N\}$ and $P'_{N+1} := \{\rho_j-\omega_j+t', t'+\rho_{i_{N+1}}\}-t-x_N = \{\rho_j-\omega_j-x_{N+1}, \rho_{i_{N+1}}-x_{N+1}\}$ are eventually coincident at s for a dense set of s in the intersection of their supports (because $T \approx_s T'$). But also, P_N and P'_N are eventually coincident at all $s \in (-\epsilon, 0) \subset [-\epsilon - x_N, \omega_{i_N} - x_N]$ as are P_{N+1} and P'_{N+1} at all $s \in (-\epsilon, 0) \subset [-\epsilon - x_{N+1}, \omega_{i_{N+1}} - x_{N+1}]$. Thus, recalling that dense eventual coincidence implies open dense eventual coincidence, we have that P_{N+1} and P_1 are eventually coincident at s for a dense set of in $[-\epsilon, 0]$, and the lemma is established by induction. □

Given a patch Q with support $[a, b]$, let $\bar{Q}$ be the periodic tiling $\bar{Q} := \cup_{k\in\mathbb{Z}}(Q+k(b-a))$. The main technical tool that we use in this article is the following theorem (which has analogues in all dimensions - see [BSW], Theorem 3.1).

Theorem 3.3. *Suppose that ϕ is a primitive Pisot substitution and suppose there is a patch Q for ϕ with support $[a, b]$ and a set $W \subset \mathbb{R}$ with the properties: the subgroup of $\mathbb{R}/(b-a)\mathbb{Z}$ generated by $\{w + (b-a)\mathbb{Z} : w \in W\}$ is dense in $\mathbb{R}/(b-a)\mathbb{Z}$; and $\bar{Q}$ and*

$\bar{Q} - w$ are densely eventually coincident for all $w \in W$. Then $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum.

Proof. Given tilings T, T' made of tiles for ϕ (but not necessarily in Ω_ϕ) let us write $T \sim T'$ provided T and T' are eventually coincident at 0. For $v \in \mathbb{R}$, let $U_v := \{x \in \mathbb{R} : \bar{Q} - x \sim \bar{Q} - v - x\}$; U_v is open and U_w is open and dense in $\mathbb{R}$ for all $w \in W$. So if $w, w' \in W$ then $U_w \cap U_{w'}$ is also open and dense in $\mathbb{R}$. For $x \in U_w \cap U_{w'}$ we have $\bar{Q} - w - x \sim \bar{Q} - x \sim \bar{Q} - w' - x$ and hence $\bar{Q} - w$ and $\bar{Q} - w'$ are densely eventually coincident. This means that $\bar{Q} - (w - w')$ and $\bar{Q}$ are densely eventually coincident. Note also that $U_v = \mathbb{R}$ for $v \in (b-a)\mathbb{Z}$. Thus, replacing W by the subgroup of $\mathbb{R}$ generated by $W \cup (b-a)\mathbb{Z}$, we may assume that W is a dense subgroup of $\mathbb{R}$.

We claim that if U_v is nonempty for some $v \in \mathbb{R}$, then U_v is dense in $\mathbb{R}$. To see this, suppose that $x \in U_v$ and fix $w \in W$. There is then $x' \in U_v \cap U_w \cap (U_w - v)$, x' arbitrarily close to x. Then $\bar{Q} - v - x' \sim \bar{Q} - x'$, $\bar{Q} - w - x' \sim \bar{Q} - x'$, and $\bar{Q} - w - (x' + v) \sim \bar{Q} - (x' + v)$. It follows that $\bar{Q} - v - (x' + w) \sim \bar{Q} - (x' + w)$; i.e., $x' + w \in U_v$. Since x' is as close to x as we wish, this shows that $x + W \subset cl(U_v)$, and thus U_v is dense in $\mathbb{R}$, as claimed.

Suppose now that $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum. There are $k \in \mathbb{N}$ and $T, T' \in \Omega_\phi$ so that $\Phi^k(T) = T$, $\Phi^k(T') = T'$, $T \sim_{srp} T'$, and $T \cap T' = \emptyset$. There are then $S, S' \in \Omega_\phi$ and $t \in \mathbb{R}$ so that $B_0[S] = B_0[T]$, $B_0[S'] = B_0[T']$, and $B_0[S - t] = B_0[S' - t]$.

Let $P \subset S$ and $P' \subset S'$ be patches with supports containing $[-1, t+1]$. We may assume that $P - s$ and $P' - s'$ are sub patches of Q for some $s, s' \in \mathbb{R}$ (otherwise, replace Q by $\Phi^m(Q)$ for sufficiently large m, so that, by primitivity, translates of P and P' occur in $\Phi^m(Q)$, and replace W by $\lambda^m W$). From $S \cap S' \neq \emptyset$ we have $(\bar{Q} - (s' - s)) \cap \bar{Q} \neq \emptyset$; so $U_{s'-s} \neq \emptyset$. By the above claim, $U_{s'-s}$ is dense in $\mathbb{R}$. Since $B_0[T'] - s' \subset P' - s' \subset \bar{Q}$ and $B_0[T] - s' \subset P - s' \subset \bar{Q} - (s' - s)$, there is $x \in spt(B_0[T'] - s') \cap spt(B_0[T] - s')$ with $\bar{Q}$ and $\bar{Q} - (s' - s)$ eventually coincident at x. Then T and T' are eventually coincident at $s' + x$. In particular, there is $n \in \mathbb{N}$ with $\Phi^n(T) \cap \Phi^n(T') \neq \emptyset$. Then $\Phi^{nk}(T) \cap \Phi^{nk}(T') \neq \emptyset$. But $\Phi^{nk}(T) = T$ and $\Phi^{nk}(T') = T'$ and $T \cap T' = \emptyset$. Thus $(\Omega_\phi, \mathbb{R})$ must have pure discrete spectrum. □

Corollary 3.4. *Suppose that ϕ is a primitive Pisot substitution and suppose there is a tile τ for ϕ and a sequence $t_n \to 0$, $t_n \neq 0$, so that for each n the set of x so that τ and $\tau - t_n$ are coincident at x is dense in $spt(\tau) \cap spt(\tau - t_n)$. Then $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum.*

Proof. Let a be the type of τ and let w be a word in the language of ϕ of the form $w = bub$, $b \in \mathcal{A}$. There is $m \in \mathbb{N}$ so that w occurs in $\phi^m(a)$ and then there is a sub patch Q of $\Phi^m(\tau)$ that follows the pattern of the word bu. Let $N \in \mathbb{N}$ be such that $\lambda^m |t_n| < \omega_b$ for all $n \geq N$. The tilings $\bar{Q}$ and $\bar{Q} - \lambda^m t_n$ are densely eventually coincident for $n \geq N$. Theorem 3.3 applies with $W = \{\lambda^m t_n : n \geq N\}$. □

The following is Lemma 10 of [B] with slightly weaker hypotheses.

Lemma 3.5. *Suppose that ϕ is Pisot, primitive, and constant on final letters and suppose that $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum. There is then $B < \infty$ so that if $(x_n, i_n)R(x_{n+1}, i_{n+1})$ for $n = 1, \dots N-1$, then $\sharp\{(x_n, i_n) : n = 1, \dots, N\} \leq B$.*

Proof. Let $\epsilon > 0$ be as in Lemma 3.2. If there is no such B then, using Lemma 3.2, there are $i \in \mathcal{A}$ and $y_n \to y \in \mathbb{R}$, $y_n \neq y$, so that the patches $P = \{\rho_1 - \omega_1, \rho_1\}$ and $P_n = \{\rho_1 - \omega_1 - y_n, \rho_i - y_n\}$ are eventually coincident at a dense set of points in an interval $I_n \subset spt(P_1) \cap spt(P_n)$ of length ϵ. There is then a subsequence and an interval $J \subset I_{n_k}$ of length $\epsilon/3$ such that either $J \subset spt(P_1 - \omega_1) \cap spt(P_1 - \omega_1 - y_{n_k})$ for all k or $J \subset spt(\rho_1) \cap spt(\rho_i - y_k)$ for all k. In either case, we have a tile τ, a sequence $x_k \to 0$, $x_k \neq 0$, and a nontrivial interval $J \subset spt(\tau) \cap spt(\tau - x_k)$ so that, for each k, τ and $\tau - x_k$ are eventually coincident at a dense subset of J. If $m \in \mathbb{N}$ is large enough so that λ^m times the length of J is larger than the length of the largest prototile then there is a tile $\tau' \in \Phi^m(\tau)$ so that τ' and $\tau' - \lambda^m x_k$ are eventually coincident at a dense set of points in $spt(\tau') \cap spt(\tau' - \lambda^m x_k)$. But then $(\Omega_\phi, \mathbb{R})$ would have pure discrete spectrum by Corollary 3.4. □

Define $\bar{R}$ on $\cup_{i=1}^d \rho_i$ to be the transitive closure of R: $(x, i)\bar{R}(y, j)$ if and only if there are (x_n, i_n), $n = 1, \dots, N$ with $(x_n, i_n)R(x_{n+1}, i_{n+1})$ for $n = 1, \dots N-1$ and $(x, i) = (x_1, i_1), (y, j) = (x_N, i_N)$.

Lemma 3.6. *Suppose that ϕ is Pisot, primitive, and constant on final letters and suppose that $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum. Then $\bar{R}$ is a closed equivalence relation.*

Proof. Clearly $\bar{R}$ is an equivalence relation. That R is closed follows from $\approx_s$ being closed; by Lemma 3.5, $\bar{R}$ is also closed. □

Under the hypotheses of the two previous lemmas, let $E_i := \{(0,i),(\omega_i,i)\}$, $E := \cup_{i=1}^d E_i$, $[E] := \{(y,j) : (y,j)\bar{R}(x,i)$ for some $(x,i) \in E_i, i \in \{1,\dots,d\}\}$, and $[E]_i := [E] \cap \rho_i$. The first coordinates of elements of $[E]_i$ partition the support of ρ_i into finitely many subintervals $I_i^1 \le I_i^2 \le \cdots \le I_i^{m(i)}$.

Lemma 3.7. *Suppose that ϕ is Pisot, primitive, and constant on final letters and suppose that $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum. If $x \in \mathring{I}_i^j$, $y \in \mathring{I}_k^n$, and $(x,i)\bar{R}(y,k)$, then $I_i^j - x = I_k^n - y$ and $(x',i)\bar{R}(y',k)$ for all $x' \in \mathring{I}_i^j$ and $y' = x' - x + y \in \mathring{I}_k^n$.*

Proof. Let $[(x,i)]$ denote the $\bar{R}$ equivalence class of (x,i) and suppose that $(x,i) \notin [E]_i$. For each $(y,j) \in [(x,i)]$ there is $t_{(x,i)}(y,j) \in \mathbb{R}$ so that $t_{(x,i)}(y,j) + y = x$. Then $t_{(x,i)}(y,j) < x < t_{(x,i)}(y,j) + \omega_j$. Let $t_{(x,i)}^{min} := max_{(y,j)\in[(x,i)]} t_{(x,i)}(y,j)$ and $t_{(x,i)}^{max} := min_{(y,j)\in[(x,i)]} t_{(x,i)}$. Note that for $(y,j),(z,k) \in [(x,i)]$ we have $t_{(x,i)}(z,k) + t_{(z,k)}(y,j) = t_{(x,i)}(y,j)$ so that $t_{(y,j)}^{min} = t_{(x,i)}^{min} + t_{(x,y)}(y,j)$, and $t_{(y,j)}^{min} = t_{(x,i)}^{min} + t_{(x,y)}(y,j)$. It follows that

for any t with $t^{min}_{(x,i)} < x-t < t^{max}_{(x,i)}$ and any $(y,j) \in [(x,i)]$ we have $(x-t,i)\bar{R}(y-t,j)$ and $t^{min}_{(y,i)} < y-t < t^{max}_{(y,i)}$. (This is clearly true if $(x,i)R(y,j)$; then apply this in finitely many steps.) If $x \in \mathring{I}^j_i$, $y \in \mathring{I}^n_k$ and $(x,i)R(y,j)$, then $I^j_i = [t^{min}_{(x,i)}, t^{max}_{(x,i)}]$, $I^n_k = [t^{min}_{(y,j)}, t^{max}_{(y,j)}]$, and we have: $I^j_i - x = I^n_k - y$ and $(x',i)\bar{R}(y',k)$ for all $x' \in \mathring{I}^j_i$ and $y' = x' - x + y \in \mathring{I}^n_k$. $\square$

Let us say that intervals I^j_i and I^n_k as in the Lemma above are equivalent and let $\{J_1, \dots, J_m\}$ be the distinct equivalence classes of the I^j_i. For each $k \in \{1, \dots, m\}$, let l_k be the common length of the $I^j_i \in J_k$. We make new prototiles $\alpha_k := ([0, l_k], k)$ and a new substitution ϕ_s on $\{1, \dots, m\}$ as follows. Let $X_\phi := \cup^d_{i=1}\rho_i/E$, let $f_\phi : X_\phi \to X_\phi$ be the map that locally stretches length by a factor of λ and follows the pattern of ϕ, and let $\pi : \cup^d_{i=1}\rho_i \to X_\phi$ be the quotient map. Note that if $\pi_1 : \Omega_\phi \to X_\phi$ is given by $\pi_1(T) = \pi((x,i))$ provided $\rho_i - x \in T$ and $x \in [0, \omega_i]$, then $f_\phi \circ \pi_1 = \pi_1 \circ \Phi$. Since $\approx_s$ is invariant under Φ, $\bar{R}$, pushed forward to X_ϕ (which we continue to call $\bar{R}$) is invariant under f_ϕ. It follows from Lemma 3.7 that $X_\phi/\bar{R}$ is also a wedge of circles, one for each J_k, and f_ϕ induces a continuous map f_{ϕ_s} on $X_{\phi_s} := X_\phi/\bar{R}$ satisfying: If $\pi_s : X_\phi \to X_{\phi_s}$ is the quotient map, then $\pi_s \circ f_\phi = f_{\phi_s} \circ \pi_s$. Now let ϕ_s be the substitution on the alphabet $\{1, \dots, m\}$ with the property that $\phi_s(k) = k_1 k_2 \cdots k_r$ if f_{ϕ_s} maps $\pi_s(\alpha_k)$ first around $\pi_s(\alpha_{k_1})$, then around $\pi_s(\alpha_{k_2})$, ..., finally around $\pi_s(\alpha_{k_r})$. Let $\alpha : \{1, \dots, d\} \to \{1, \dots, m\}$ be the morphism $\alpha(i) := i_1 \cdots i_{m(i)}$ with i_j defined by $I^j_i \in J_{i_j}$; we have $\alpha \circ \phi = \phi_s \circ \alpha$.

Lemma 3.8. *Suppose that ϕ is primitive, Pisot, and constant on final letters and suppose that $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum. Then ϕ_s is primitive, Pisot, and constant on final letters and if ϕ is injective on initial letters, so is ϕ_s.*

Proof. If J_k is an equivalence class of intervals, each $I_i^j \in J_k$ has the property that its endpoints $a < b$ satisfy (a, i) and (b, i) are each $\bar{R}$-equivalent to an endpoint of some prototile: say $(a, i)\bar{R}(0, r)$ and $(b, i)\bar{R}(\omega_t, t)$. Then $I_r^1, I_t^{m(t)} \in J_k$. Let $l \in \mathcal{A}$ be such that $\phi(c) = \cdots l$ for each $c \in \mathcal{A}$ and let l' be such that $I_l^{m(l)} \in J_{l'}$. Then

$$\phi_s \circ \alpha(t) = \phi_s(\cdots k) = \cdots \phi_s(k).$$

On the other hand,

$$\alpha \circ \phi(t) = \alpha(\cdots l) = \cdots l'.$$

From $\phi_s \circ \alpha = \alpha \circ \phi$, we have $\phi_s(k) = \cdots l'$ for all k; that is, ϕ_s is constant on final letters.

Primitivity of ϕ_s follows from primitivity of ϕ, the relation $\alpha \circ \phi = \phi_s \circ \alpha$, and surjectivity of π_s.

If ϕ is injective on initial letters there is $n \in \mathbb{N}$ so that $\phi^n(c) = c \cdots$ for all $c \in \mathcal{A}$. Then from $\alpha(k) = r \cdots$ we have

$$\phi_s^n \circ \alpha(r) = \phi_s^n(k \cdots) = \phi_s^n(k) \cdots$$

and also

$$\alpha \circ \phi^n(r) = \alpha(r \cdots) = k \cdots ,$$

from which it follows that $\phi_s^n(k) = k \cdots$ for all k, and ϕ_s is injective on initial letters.

□

Theorem 3.9. *Suppose that ϕ is primitive, Pisot, and constant on final letters and suppose that $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum. Then the substitution ϕ_s is Pisot and the dynamical system $(\Omega_\phi/\approx_s, \mathbb{R})$ is topologically isomorphic with $(\Omega_{\phi_s}, \mathbb{R})$ by an isomorphism that conjugates the substitution homeomorphisms. Furthermore, the relation $\approx_s$ is trivial on Ω_{ϕ_s} and $(\Omega_{\phi_s}, \mathbb{R})$ does not have pure discrete spectrum.*

Proof. This is basically the 1-dimensional of Theorem 12 of [B]. We provide an argument here for the reader's convenience.

The morphism α induces a map $\bar{\alpha} : \Omega_\phi \to \Omega_{\phi_s}$ that semi conjugates Φ with Φ_s and also the $\mathbb{R}$-actions on the two spaces (see, for example, [BD2]). We will show that $\bar{\alpha}(T) = \bar{\alpha}(T')$ if and only if $T \approx_s T'$.

Let $\pi_1 : \Omega_\phi \to X_\phi$ be given by $\pi_1(T) = [(x, i)]$ if and only if $\rho_i - x \in T$ and, likewise, let $\pi_2 : \Omega_{\phi_s} \to X_{\phi_s}$ be defined by $\pi_2(S) = [(x, k)]$ if and only if $\alpha_k - x \in S$. We have

the following commuting diagram.

$$\begin{array}{ccc} \Phi \circlearrowleft \Omega_\phi & \xrightarrow{\bar{\alpha}} & \Omega_{\phi_s} \circlearrowleft \Phi_s \\ \downarrow \pi_1 & & \downarrow \pi_2 \\ f_\phi \circlearrowleft X_\phi & \xrightarrow{\pi_s} & X_{\phi_s} \circlearrowleft f_{\phi_s} \end{array} \tag{1}$$

Now if $T \approx_s T'$ (in Ω_ϕ), then $T - t \approx_s T' - t$ for all $t \in \mathbb{R}$, so $\pi_1(T-t)\bar{R}\pi_1(T'-t)$, i.e., $\pi_s(\pi_1(T-t)) = \pi_s(\pi_1(T'-t))$, and hence $\pi_2(\bar{\alpha}(T)-t) = \pi_2(\bar{\alpha}(T')-t)$, for all $t \in \mathbb{R}$. This means that $\bar{\alpha}(T) = \bar{\alpha}(T')$.

Conversely, suppose that $\bar{\alpha}(T) = \bar{\alpha}(T')$. Then $\bar{\alpha}(T-t) = \bar{\alpha}(T'-t)$ for all $t \in \mathbb{R}$ and from diagram 1 we have $\pi_1(T-t)\bar{R}\pi_1(T'-t)$ for all $t \in \mathbb{R}$. The set U of $t \in \mathbb{R}$ so that neither $\pi_1(T-t)$ nor $\pi_1(T'-t)$ equals the branch point E of X_ϕ is open and dense in $\mathbb{R}$. Fix $t \in U$, let $[(x,i)] = \pi_1(T-t)$, and let $[(x',j)] = \pi_1(T'-t)$. Then $(x,i)\bar{R}(x',j)$ and, from Lemma 3.2, there is $\epsilon > 0$ so that the patches $\{\rho_1 - \omega_1 - x, \rho_i - x\}$ and $\rho_1 - \omega_1 - x', \rho_j - x'\}$ are densely eventually coincident at s for a dense set of s in $[-\epsilon, 0]$. This means that $T-t$ and $T'-t$ are densely eventually coincident at a dense set of s in $[-\epsilon', 0]$ where $\epsilon' > 0$ is such that $[-\epsilon, 0] \subset spt(\rho_i - x) \cap spt(\rho_j - x')$. It follows that T and T' are densely eventually coincident. It remains to show that T and T' are strongly regionally proximal.

From Lemma 3.2 and Corollary 3.4 it follows that there are, up to translation, only finitely many pairs $\{\rho_i - x, \rho_j - x'\}$ with $(x,i)\bar{R}(x',j)$. We are assuming that $\bar{\alpha}(T) = \bar{\alpha}(T')$. From diagram 1, $\bar{\alpha}(\Phi^{-k}(T)) = \bar{\alpha}(\Phi^{-k}(T'))$ for all $k \in \mathbb{N}$. There

are then $i, j \in \mathcal{A}, x_n \geq 0, x'_n \geq 0$, and $k_n \to \infty$ so that $\rho_i - x_n \in B_0[\Phi^{-k_n}(T)]$, $\rho_j - x'_n \in B_0[\Phi^{-k_n}(T')]$, and $x_n - x'_n$ is constant. Since $\Phi^{-k_1}(T)$ and $\Phi^{-k_1}(T')$ are densely eventually coincident, there is $s \in spt(\rho_i - x_1) \cap spt(\rho_j - x'_1)$, $N \in \mathbb{N}$, and a tile η so that $\eta \in \Phi^N(\rho_i - x_1) \cap \Phi^N(\rho_j - x'_1)$. Then, for $k_n \geq N$, $\Phi^{k_n - N}(\eta) + \lambda^{k_n - N}(x_N - x_1) \in T \cap T'$. Thus T and T' are strongly proximal, and hence strongly regionally proximal. We have shown that $\bar{\alpha}(T) = \bar{\alpha}(T') \Leftrightarrow T \approx_s T'$. Since the $\mathbb{R}$-action on Ω_{ϕ_s} is minimal and $\bar{\alpha}(T - t) = \bar{\alpha}(T) - t$, $\bar{\alpha}$ is surjective, and hence induces an isomorphism of $(\Omega_\phi / \approx_s, \mathbb{R})$ with $(\Omega_{\phi_s}, \mathbb{R})$.

To see that $\approx_s$ is trivial on $(\Omega_{\phi_s}, \mathbb{R})$, suppose that $\bar{\alpha}(T) \approx_s \bar{\alpha}(T')$. In particular, there is a dense set U so that for each $t \in U$ there is $n(t) \in \mathbb{N}$ with $B_0[\Phi_s^{n(t)}(\bar{\alpha}(T) - t)] = B_0[\Phi_s^{n(t)}(\bar{\alpha}(T') - t)]$. Fix such a t. Then $\pi_2(\Phi_s^{n(t)}(\bar{\alpha}(T) - t)) = \pi_2(\Phi_s^{n(t)}(\bar{\alpha}(T') - t))$, so $(\Phi^{n(t)}(T) - \lambda^{n(t)}t) \bar{R} (\Phi^{n(t)}(T') - \lambda^{n(t)}t)$, by diagram 1. Using Lemma 3.2 (as above), there is then $\epsilon' > 0$ so that $\Phi^{n(t)}(T) - \lambda^{n(t)}t$ and $\Phi^{n(t)}(T') - \lambda^{n(t)}t$ are eventually coincident at a dense set of points in $[-\epsilon', 0]$. Then $T - t$ and $T' - t$ are eventually coincident at a dense set of points in $[-\epsilon' \lambda^{-n(t)}, 0]$ and, since the set of such t is dense, T and T' are densely eventually coincident.

We now argue that if $\bar{\alpha}(T) \approx_s \bar{\alpha}(T')$, then T and T' are strongly regionally proximal (denoted $T \sim_{srp} T'$). Let $\pi_{max} : \Omega_\phi \to \Omega_{max}$ and $\pi^s_{max} : \Omega_{\phi_s} \to \Omega^s_{max}$ denote the maximal equicontinuous factor maps for the systems $(\Omega_\phi, \mathbb{R})$ and $(\Omega_{\phi_s}, \mathbb{R})$, resp., and recall

that tilings T, T' in either space have the same image under the maximal equicontinuous factor map if and only if $T \sim_{srp} T'$. Since $\bar{\alpha}(T) = \bar{\alpha}(T')$ implies that T and T' are strongly regionally proximal, π_{max} factors through Ω_{ϕ_s}. Hence, by maximality, and by uniqueness up to isomorphism of the maximal equicontinuous factor, $(\Omega_{max}, \mathbb{R})$ and $(\Omega^s_{max}, \mathbb{R})$ are isomorphic and we may take $\Omega^s_{max} = \Omega_{max}$ and $\pi_{max} = \pi^s_{max} \circ \bar{\alpha}$. Now $\bar{\alpha}(T) \approx_s \bar{\alpha}(T') \implies \bar{\alpha}(T) \sim_{srp} \bar{\alpha}(T') \implies \pi^s_{max}(\bar{\alpha}(T)) = \pi^s_{max}(\bar{\alpha}(T')) \implies T \sim_{srp} T'$. Together with $\bar{\alpha}(T) \approx_s \bar{\alpha}(T')$ implies T and T' are densely eventually coincident, this shows that $\bar{\alpha}(T) \approx_s \bar{\alpha}(T') \implies T \approx_s T' \implies \bar{\alpha}(T) = \bar{\alpha}(T')$; i.e., $\approx_s$ is trivial on Ω_{ϕ_s}.

For each $z \in \Omega_{max} = \Omega^s_{max}$ there are $T, T' \in \pi^{-1}_{max}(z)$ with T and T' nowhere eventually coincident ([BK]). Then $\bar{\alpha}(T) \neq \bar{\alpha}(T')$ but $\pi^s_{max}(\bar{\alpha}(T)) = \pi^s_{max}(\bar{\alpha}(T'))$. Thus, π^s_{max} is at least 2-1 everywhere and $(\Omega_{\phi_s}, \mathbb{R})$ does not have pure discrete spectrum.

□

Lemma 3.10. *Suppose that ϕ is primitive, non-periodic, Pisot, injective on initial letters, and constant on final letters. Then there are $i \neq j \in \mathcal{A}$ so that the prototiles ρ_i and ρ_j are eventually coincident at a dense set of points in $spt(\rho_i) \cap spt(\rho_j)$.*

Proof. By replacing ϕ by ϕ^k for an appropriate $k \in \mathbb{N}$, we may assume that $\phi(a) = a \cdots$ for all $a \in \mathcal{A}$. Let r be the coincidence rank of ϕ. There are then $T_1, \ldots, T_r \in \Omega_\phi$ with the properties: $T_i \sim_{srp} T_j$ for all $i, j \in \{1, \ldots, r\}$; $\Phi^n(T_i) \cap \Phi^n(T_j) = \emptyset$ for all $i \neq j \in \{1, \ldots, r\}$ and all $n \in \mathbb{N}$; and the T_i are Φ-periodic (see [BK], or Theorem 4

of [B]). Replacing ϕ by an appropriate power, we may assume that the T_i are all fixed by Φ. It follows from the constancy of ϕ on final letters that, for $i \neq j \in \{1, \dots, r\}$, the set of endpoints of T_i (that is, the set of endpoints of supports of tiles in T_i) is disjoint from the set of endpoints of T_j. Let $V = \{\cdots < v_{-1} < v_0 < v_1 < \cdots\}$ denote the union of the sets of endpoints of the T_i, $i = 1, \dots, r$. For each $k \in \mathbb{Z}$ and $i \in \{1, \dots, r\}$ let τ_i^k denote the tile in T_i that has $(v_k + v_{k+1})/2$ in its support. We will call $\mathcal{C}_k := \{\tau_1^k, \dots, \tau_r^k\}$ a *configuration*. Up to translation, there are only finitely many distinct configurations (see [BK]), and since the T_i are not (translation) periodic, the translation equivalence of $\mathcal{C}_k$ is not a periodic function of k. Therefore, there are $k, l \in \mathbb{Z}$ so that $\mathcal{C}_{k-1} - v_{v_k - 1} = \mathcal{C}_{l-1} - v_{l-1}$ while $\mathcal{C}_k - v_k \neq \mathcal{C}_l - v_l$. For any $n \in \mathbb{Z}$, the configurations $\mathcal{C}_{n-1}$ and $\mathcal{C}_n$ differ only in the tile of $\mathcal{C}_{n-1}$ with terminal endpoint v_n and the tile of $\mathcal{C}_n$ with initial endpoint v_n. Hence there are $i \neq j$ with $\rho_i \in \mathcal{C}_k - v_k$, $\rho_j \in \mathcal{C}_l - v_l$, and $(\mathcal{C}_k - v_k) \setminus \{\rho_i\} = (\mathcal{C}_l - v_l) \setminus \{\rho_j\}$.

Suppose (by way of contradiction) that ρ_i and ρ_j are not eventually coincident at a dense set of points in the intersection of their supports. Note that because $\phi(i) = i \cdots$ and $\phi(j) = j \cdots$, eventual coincidence of ρ_i and ρ_j at a dense set of points in an interval $[0, \delta]$, with $\delta > 0$, would imply eventual coincidence of those prototiles at a dense set of points in the intersection of their supports. Thus there are $0 \leq t_0 < t_2 \leq min\{v_{k+1} - v_k, v_{l+1} - v_l\}$ so that ρ_i and ρ_j are not eventually coincident at any point of

the interval $[t_0, t_2]$. Let i', j' be such that $\rho_i \in T_{i'} - v_k$ and $\rho_j \in T_{j'} - v_l$ and let $t_1 := \frac{t_0+t_2}{2}$. By compactness, there are $n_p \to \infty$ so that $\Phi^{n_p}(T_q - v_k - t_1) \to S_q \in \Omega_\phi$, for all $q \in \{1, \ldots, r\}$, and $\Phi^{n_p}(T_{j'} - v_l - t_1) \to S_{r+1} \in \Omega_\phi$. We claim that the S_q, $q = 1, \ldots, r+1$, are pairwise strongly regionally proximal. Indeed, strong regional proximality is closed and Φ-invariant, so the S_q, $q = 1, \ldots, r$, are strongly regionally proximal. Pick $q' \in \{1, \ldots, r\} \setminus \{j_1\}$. There is then $q'' \in \{1, \ldots, r\}$ with $B_0[T_{q'} - v_l - t_1] = B_0[T_{q''} - v_k - t_1]$. Then $\Phi^{n_p}(T_{q'} - v_l - t_1) \to S_{q''}$, and again, since $\sim_{srp}$ is closed and Φ-invariant, S_{r+1} is strongly regionally proximal with $S_{q''}$. Hence the S_q, $q = 1, \ldots, r+1$, are all strongly regionlly proximal, as claimed. We further claim that the S_q, $q \in \{1, \ldots, r+1\}$, are pairwise disjoint. For $q \in \{1, \ldots, r\}$ this is clear: since the T_q are fixed by Φ, the set of configurations formed by the $\Phi^{n_p}(T_q - v_k - t_1)$ equals the set of configurations formed by the T_q and hence the set of configurations formed by the S_q is contained in the set of configurations formed by the T_q. In particular, each configuration formed by the S_q, $q \in \{1, \ldots, r\}$, consists of r distinct tiles, so these S_q are pairwise disjoint. The same argument shows that S_{r+1} is disjoint from S_q for $q \in \{1, \ldots, r\} \setminus \{i_1\}$, since $\{S_q : q \in \{1, \ldots, r+1\} \setminus \{i_1\}\} = \{\lim_{p\to\infty} \Phi^{n_p}(T_q - v_l - t_1) : q \in \{1, \ldots, r\}\}$. It remains to show that $S_{r+1} \cap S_{i_1} = \emptyset$. For this, consider the tilings $U_i, U_j \in \Omega_\phi$ that are fixed by Φ with $\rho_c - \omega_c, \rho_i \in U_i$ and $\rho_c - \omega_c, \rho_j \in U_j$, where c is such that $\phi(a) = \cdots c$ for all $a \in \mathcal{A}$. Then $S_{i_1} = \lim_{p\to\infty} \Phi^{n_p}(U_i - t_1)$, and $S_{r+1} = \lim_{p\to\infty} \Phi^{n_p}(U_j -$

t_1). As $U_i \sim_{srp} U_j$ (since U_i and U_j are proximal), there are, up to translation, only finitely many pairs $\{B_0[U_i - t], B_0[U_j - t]\}$, $t \in \mathbb{R}$. It follows that if there are t_p with $B_0[\lim_{p\to\infty}(U_i - t_p)] = B_0[\lim_{p\to\infty}(U_j - t_p)]$, then $B_0[U_i - t_p] = B_0[U_j - t_p]$ for all large p. Thus, if $S_{i_1} \cap S_{r+1} \neq \emptyset$, say $B_0[S_{i_1} - t] = B_0[S_{r+1} - t]$, and $t_p = t_1\lambda^{n_p} - t$, we have $B_0[U_i - t_p] = B_0[\Phi^{n_p}(U_i - t_1) - t] = B_0[\Phi^{n_p}(U_j - t_1) - t] = B_0[U_j - t_p]$ for all large p. For p large enough that $|t/\lambda^{n_p}| < (t_0 + t_2)/2$, we have $s := t_1 - (t/\lambda^{n_p}) \in (t_0, t_2)$ with ρ_i and ρ_j eventually coincident at s, a contradiction. This proves that S_{i_1} and S_{r+1} are disjoint and hence the claim that the S_q, $q \in \{1, \dots, r+1\}$, are pairwise disjoint.

We now have the situation that there are $r+1$ tilings $S_1, \dots, S_{r+1}$ that are strongly regionally proximal and pairwise disjoint. Suppose that $T' = T'_1 \in \Omega_\phi$. By minimality of the $\mathbb{R}$-action, there are s_n such that $S_1 - s_n \to T'_1$, by passing to a subsequence, we may assume that $S_q - s_n \to T'_q \in \Omega_\phi$ for $q = 1, \dots, r+1$. Then the T'_q are strongly regionally proximal and pairwise disjoint (the latter uses again that, up to translation, there are only finitely many pairs $\{B_0[S_q - t], B_0[S_{q'} - t]\}$). But then the coincidence rank of ϕ is at least $r+1$, not r. Thus ρ_i and ρ_j must be eventually coincident at a dense set of points in the intersection of their supports. □

Corollary 3.11. *Suppose that ϕ is primitive, non-periodic, Pisot, injective on initial letters, and constant on final letters. Then the relation $\approx_s$ is nontrivial on Ω_ϕ.*

Proof. Say $\phi(a) = \cdots c$ for all $a \in \mathcal{A}$. Let ρ_i, ρ_j be as in Lemma 3.10 and let $U_i, U_j \in \Omega_\phi$ be the Φ-periodic tilings with $\rho_c - \omega_c, \rho_i \in U_i$ and $\rho_c - \omega_c, \rho_j \in U_j$. Then $U_i \neq U_j$ and $U_i \approx_s U_j$. □

Theorem 3.12. *Suppose that ϕ is primitive, Pisot, injective on initial letters, and constant on final letters. Then $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum.*

Proof. Suppose that $(\Omega_\phi, \mathbb{R})$ does not have pure discrete spectrum. Then ϕ_s is also primitive, Pisot, injective on initial letters, and constant on final letters by Lemma 3.8. By Theorem 3.9, $\approx_s$ is trivial on Ω_{ϕ_s}. If there were a translation periodic tiling in Ω_{ϕ_s}, it would follow from primitivity of ϕ_s that $(\Omega_{\phi_s}, \mathbb{R})$, being simply translation on a circle, has pure discrete spectrum, which it doesn't by Theorem 3.9. Thus ϕ_s is non-periodic. But now Corollary 3.11 says $\approx_s$ is nontrivial on Ω_{ϕ_s}. Thus $(\Omega_\phi, \mathbb{R})$ must have pure discrete spectrum. □

Remark 3.13. *By replacing ϕ by ϕ^n for appropriate $n \in \mathbb{N}$, the hypothesis in Theorem 3.12 that ϕ is constant on final letters can be weakened to ϕ being eventually constant on final letters. Also, if ϕ satisfies the other hypotheses but is not necessarily primitive, let c be the (eventual) last letter of all $\phi(a)$ and let $\mathcal{A}' := \{a \in \mathcal{A} : \exists n \in \mathbb{N}$ with a occurring in $\phi^n(c)\}$. Then the theorem applies to $\phi' := \phi|_{\mathcal{A}'}$ to conclude that the 'minimal core' $(\Omega_{\phi'}, \mathbb{R})$ of $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum.*

4. Applications

4.1. **β-substitutions.** Given $\beta > 1$, let $T_\beta : [0,1] \to [0,1]$ be the *β-transformation* defined by $x \mapsto \beta x - \lfloor \beta x \rfloor$. The number β is called a *Parry number* if the orbit of 1 under T_β is finite and a *simple Parry number* if $T_\beta^n(1) = 0$ for some $n \in \mathbb{N}$. All Pisot numbers are Parry numbers ([Ber, Sc]).

Given a simple Parry number β with n (smallest) as above, let $P_i := [0, T_\beta^{i-1}(1)]$ for $i = 1, \dots, n$. Then, for $i = 1, \dots, n-1$, T_β maps P_i a_i times across $P_1 = [0,1]$ and once across P_{i+1}, and T_β maps P_n exactly a_n times across P_1. The *β-substitution*, ϕ_β, is given by:

$$\begin{aligned} \phi_\beta(1) &= 21^{a_1} \\ \phi_\beta(2) &= 31^{a_2} \\ &\vdots \\ \phi_\beta(n) &= 1^{a_n}. \end{aligned}$$

Taking $\rho_i := (P_i, i)$, $i = 1, \dots, n$, as prototiles for ϕ_β, we define (a.e.) the map $p : \Omega_{\phi_\beta} \to \varprojlim T_\beta$ by $p(T) := (\dots, t_{-1}, t_0, t_1, \dots)$ with t_k determined by: $\rho_i - t_k \in \Phi_\beta^{-k}(T)$ and $t_k \in int(spt(\rho_i))$ for some i. Then p is a metric isomorphism that conjugates the

substitution-induced homeomorphism on Ω_{ϕ_β} with the shift homeomorphism $\hat{T}_\beta$ on the inverse limit space $\varprojlim T_\beta$.

Each non-negative real number x has a *greedy expansion* in base β:

$$x = \sum_{k=-N}^{\infty} x_k \beta^{-k}$$

with $x_k \in \{0, \ldots, \lfloor \beta \rfloor\}$ satisfying $|x_M - \sum_{k=-N}^{M} x_k \beta^{-k}| < \beta^{-M}$ for each $M \geq -N$. Each such greedy expansion determines $(\ldots, 0, 0, x_{-N}, \ldots, x_{-1}, x_0, x_1, \ldots) \in \{0, \ldots, \lfloor \beta \rfloor\}^{\mathbb{Z}}$; the closure of all such sequences is denoted by Σ_β and called the *β-shift*.[1] For β a simple Parry number, Σ_β is a subshift of finite type. The map $r : (\ldots, x_{-1}, x_0, x_1, \ldots) \mapsto (\ldots, \sum_{k=1}^{\infty} x_{k-1}\beta^{-k}, \sum_{k=1}^{\infty} x_k \beta^{-k}, \sum_{k=1}^{\infty} x_{k+1}\beta^{-k}, \ldots)$ defines a metric isomorphism that conjugates the shift σ on Σ_β with $\hat{T}_\beta$ on $\varprojlim_{T_\beta}$. The map

$$g := \pi_{max} \circ p^{-1} \circ r : \Sigma_\beta \to \Omega_{max},$$

where $\pi_{max} : \Omega_{\phi_\beta} \to \Omega_{max}$ is the maximal equicontinuous factor map, gives a continuous and bounded-to-one semi-conjugacy between the β-shift σ and the hyperbolic automorphism $(\Phi_\beta)_{max} : \Omega_{max} \to \Omega_{max}$. This map g has the nice property of being an *arithmetical coding*[2]: If x and x' are non-negative real numbers with sequences $\underline{x}, \underline{x'}$ and $\underline{x + x'}$ in Σ_β corresponding to the greedy β-expansions of x, x' and $x + x'$, then

[1] Not to be confused with the substitutive system Σ_{ϕ_β}.

[2] The terminology is due to Siderov ([Si]).

$g(\underline{x+x'}) = g(\underline{x}) + g(\underline{x'})$. Since ϕ_β is primitive, injective on initial letters, and eventually constant on final letters (see Remark 3.13), it follows from Theorem 3.12 that, for Pisot simple Parry numbers, the map π_{max}, and hence the coding g, is a.e. one-to-one. (This result appears in [BBK] under the additional hypothesis that the algebraic degree of β is greater than n/p, p the smallest prime divisor of n.)

Corollary 4.1. *If β is a Pisot simple Parry number, then the system $(\Omega_{\phi_\beta}, \mathbb{R})$ has pure discrete spectrum.*

Remark 4.2. *For β a Pisot simple Parry number and a unit, the substitutive system $(\Sigma_{\phi_\beta}, \mathbb{Z})$ also has pure discrete spectrum provided the substitution ϕ_β is irreducible (that is, $n = d$, d the algebraic degree of β -* [BKw] *or* [CS]*). But if ϕ_β is reducible this is not necessarily the case as counterexamples of Ei and Ito show (*[EI]*). In any event, if β is a unit, $(\Sigma_{\phi_\beta}, \mathbb{Z})$ is measurably conjugate to the system of a map induced by a rotation on the $(d-1)$-torus (*[BBK]*, Proposition 8.1).*

4.2. **Arnoux-Rauzy, Brun, and Jacobi-Perron substitutions.** The substitutions of this section arise as finite products of elements taken from certain collections of basic substitutions. The Arnoux-Rauzy substitutions generalize the two-letter Sturmian substitutions and the three-letter Rauzy substitution (see [BFZ] and [CC] for general accounts of these) and the Brun and Jacobi-Perron substitutions come from multidimensional continued fraction algorithms (see [Be] and [S]).

Let $\mathcal{A} = \{1, \ldots, d\}$ and for each $i \in \mathcal{A}$, let σ_i be the substitution on $\mathcal{A}$ defined by $\sigma_i(i) = i$ and $\sigma_i(j) = ji$ for $j \neq i$. Given a word $w = w_1 \cdots w_k \in \mathcal{A}^*$ that contains at least one occurrence of each letter of $\mathcal{A}$, let $\sigma_w = \sigma_{w_1} \circ \cdots \circ \sigma_{w_k}$. Such a σ_w is called an *Arnoux-Rauzy substitution.* Arnoux and Ito ([AI]) first proved that all Arnoux-Rauzy substitutions on 2 or 3 letters are irreducible Pisot and recently Avila and Delecroix have proved that Arnoux-Rauzy substitutions on any number of letters are irreducible Pisot ([AD]). The following (assuming Pisot) is proved in [BSW] by analyzing balanced pairs and a rather different proof, for $d = 3$, has been given by Berthé, Jolivet, and Siegel ([BJS]). It is easily verified that Arnoux-Rauzy substitutions are primitive, injective on initial letters, and constant on final letters.

Corollary 4.3. *If σ_w is an Arnoux-Rauzy substitution, then $(\Omega_{\sigma_w}, \mathbb{R})$ has pure discrete spectrum.*

The *Brun substitutions*

$$\sigma_1^{Brun} : 1 \mapsto 1,\ 2 \mapsto 2,\ 3 \mapsto 32;$$

$$\sigma_2^{Brun} : 1 \mapsto 1,\ 2 \mapsto 3,\ 3 \mapsto 23;\ \text{and}$$

$$\sigma_3^{Brun} : 1 \mapsto 2,\ 2 \mapsto 3,\ 3 \mapsto 13$$

are discussed by Berthé, Bourdon, Jolivet, and Siegel in [BBJS] where they prove that if $w = w_1 \cdots w_k \in \{1, 2, 3\}^*$ is any word containing at least one occurrence of 3 and

$\phi = \sigma_{w_1}^{Brun} \circ \cdots \circ \sigma_{w_k}^{Brun}$, then $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum.

The *Jacobi-Perron substitutions* are

$$\sigma_{a,b}^{JP} : 1 \mapsto 3,\ 2 \mapsto 13^a,\ 3 \mapsto 23^b$$

for $a, b \in \mathbb{N}_0$. Any product $\phi = \sigma_{a_1,b_1}^{JP} \circ \cdots \circ \sigma_{a_n,b_n}^{JP}$ is irreducible Pisot as long as $0 \leq a_i \leq b_i$ and $b_i \neq 0$ for $i = 1, \ldots, n$ ([DFP]). It is proved in [BBJS] that, for such ϕ, $(\Omega_\phi, \mathbb{R})$ has pure discrete spectrum.

The results for the Brun and Jacobi-Perron substitutions follow immediately from Theorem 3.12 and Remark 3.13.

References

[ABBLS] S. Akiyama, M. Barge, V. Berthé, J.-Y. Lee, A. Siegel, On the Pisot Conjecture, 2012 preprint.

[AL1] S. Akiyama and J.-Y. Lee, Algorithm for determining pure pointedness of self-affine tilings, *Adv. Math.* **226**, no. 4, (2011), 2855-2883.

[AL2] S. Akiyama and J.-Y. Lee, Computation of pure discrete spectrum of self-affine tilings. Preprint, 2013.

[AP] J.E. Anderson and I.F. Putnam, Topological invariants for substitution tilings and their associated C^*-algebras, *Ergodic Theory & Dynamical Systems* **18** (1998), 509–537.

[AI] P. Arnoux and S. Ito, Pisot Substitutions and Rauzy fractals, *Bull. Belg. Math Soc.* **8** (2001), 181-207.

[AD] A. Avila and V. Delecroix, Pisot property for constant cocycles over shift of finite type, 2013 preprint.

[BBK] V. Baker, M. Barge and J. Kwapisz, Geometric realization and coincidence for reducible non-unimodular Pisot tiling spaces with an application to β-shifts, *Ann. Inst. Fourier (Grenoble)* **56** No. 7 (2006), p. 2213–2248.

[B] M. Barge, Factors of Pisot tiling spaces and the Coincidence Rank Conjecture, arXiv:1301.7094, (2013).

[BD2] M. Barge, B. Diamond, A complete invariant for the topology of one- dimensional substitution tiling spaces, *Ergod. Th. & Dyn. Sys.* **21** (2001), 1333-1358.

[BD1] M. Barge and B. Diamond, Coincidence for substitutions of Pisot type, *Bull. Soc. Math. France* **130** (2002), p. 691–626.

[BBJS] V. Berthé, J. Bourdon, T. Jolivet, and A. Siegel, A combinatorial approach to products of Pisot substitutions, arXiv:1401.0704v1, (2014).

[BK] M. Barge and J. Kellendonk, Proximality and pure point spectrum for tiling dynamical systems, arXiv:1108.4065, (2012), to appear in Michigan Math. J..

[BKw] M. Barge and J. Kwapisz, Geometric theory of unimodular Pisot substitutions, *Amer J. Math.* **128** (2006), 1219-1282.

[BSW] M. Barge, S. Štimac and R.F. Williams, Pure discrete spectrum in substitution tiling spaces, *Disc. and Cont. Dynam. Sys. - A***2** (2013), 579-597.

[BFZ] V. Berthé, S. Ferenczi, and L. Q. Zamboni, Interactions between dynamics, arihtmetics and combinatorics: the good, the bad, and the ugly, *Algebraic and topological dynamics, Contemp. Math.*, **385**, Amer. Math. Soc., Providence, RI (2005), 333-364.

[BJS] V. Berthé, T. Jolivet and A. Siegel, Substitutive Arnoux-Rauzy substitutions have pure discrete spectrum, *Unif. Distrib. Theory* **7**, no. 1, (2012), 173-197.

[Be] V. Berthé, Multidimensional Euclidean algorithms, numeration and substitutions, *Integers* **11B** (2011), Paper No. A2, 34.

[Ber] A. Bertrand, Développements en base de Pisot et répartition modulo 1, *C. R. Acad. Sci. Paris*, **285**, no.6, (1977), A419-A421.

[CC] J. Cassaigne and N. Chekhova, Fonctions de récurrence des suites d'Arnoux-Rauzy et réponse à une questionde Morseet Hedlund, *Ann. Inst. Fourier (Grenoble)* **56**, no. 7, Numération, pavages, substitutions (2006), 2249-2270.

[CS] A. Clark and L. Sadun, When shape matters: deformations of tiling spaces, *Ergodic Theory Dynam. Systems*, **26** (2006), 69-86.

[DFP] E. Dubois, A. Farhane, and R. Paysant-Le Roux, The Jacobi-Perron algorithm and Pisot numbers, *Acta Arith.* **111** (2004), 269-275.

[EI] H. Ei and S. Ito, Tilings from some non-irreducible, Pisot substitutions,*Discrete Math. and Theo. Comp. Science* **8**, no.1 (2005), 81-122.

[HS] M. Hollander and B. Solomyak, Two-symbol Pisot substitutions have pure discrete spectrum, *Ergodic Theory & Dynamical Systems* **23** (2003), 533-540.

[M] B. Mossé, Puissances de mots et reconnaissabilité des points fixes d'une substitution, *Theoretical Computer Science* **99** (1992), 327-334.

[S] F. Schweiger, *Multidimensional continued fractions*, Oxford Science Publications, Oxford University Press, Oxford, 2000.

[Sc] K. Schmidt, On periodic expansions of Pisot numbers and Salem numbers, *Bull. London Math. Soc.*, **12** (1980), 269-278.

[Si] N. Sidorov, Arithmetic dynamics, in *Topics in dynamics and ergodic theory*, vol. 310 of *London Math. Soc. Lecture Note Ser.*, pp. 145-189, Cambridge: Cambridge Univ. Press, 2003.

[S1] B. Solomyak, Dynamics of self-similar tilings, *Ergod. Th. & Dynam. Sys.*, **17** (1997), 695-738.

[S2] B. Solomyak, Nonperiodicity implies unique composition for self-similar translationally finite tilings, *Discrete Comput. Geometry* **20** (1998), 265–279.

[W] P. Walters, *An introduction to ergodic theory*, Springer-Verlag, New York, 1982.

Montana State University, Bozeman, MT, 59717

E-mail address: barge@math.montana.edu